IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

ATTY.'s DOCKET: MUKAI=2

In re Application of:

(Application of:

DECLARRATION UNDER 37 CFR 1.132

Honorable Commissioner of Patents and Trademarks Washington, D.C. 20231

Sir:

- 1. I, Tomoyuki NISHIMOTO, declare as follows:
- 2. I am a citizen of Japan, residing at 500-30 Meguro-cho, Okayama-shi, Okayama, Japan.
- 3. In 1985, I received a bachelor of Agriculture from Osaka Prefecture University, and in 1998 I received a doctorate of Agriculture from the above-identified university.
- 4. As shown in my curriculum vitae attached hereto as Attachment A, from 1990 to 2009, I researched in Hayashibara Biochemical Laboratories Inc. fundamental studies and industrial applications of carbohydrates and their related enzymes. Since 2009, I have been a director of Research

Center, Hayashibara Biochemical Laboratories, Inc.

- I have read and thoroughly understood the present invention and the content of the United States Patent No. 5,137,723, titled " α -GLYCOSYL-L-ASCORBIC ACID, AND ITS PREPARATION AND USES" applied for by "Kabushiki Kaisha Hayashibara Seibutsu Kagaku Kenkyujo" (Hayashibara Biochemical Laboratories, Inc.), cited in an official action in the procedure of the present invention.
- In the declaration dated November 28, 2008, I demonstrated on the 6. α -isomaltosyl the results that experimental the of basis glucosaccharide-forming enzyme (abbreviated as "IMG", hereinafter) according to the present invention is significantly different from rat intestine $\alpha\text{-glucosidase}$ (RIAGase) described in the above-identified patent in the productivity of 2-O- α -D-glucopyranosyl-L-ascorbic acid (AA2G), and (AA5G), acid $5-O-\alpha-D-glucopyranosyl-L-ascorbic$ $6\text{-}0\text{-}\alpha\text{-}D\text{-}glucopyranosyl-L-ascorbic}$ acid (AA6G). The experiment was conducted using "PINEDEX® #1", a partial starch hydrolyzate having a dextrose equivalent (DE) of about 8 ± 1 , as a substrate.
 - 7. This time, I conducted the following experiment using three kinds of partial starch hydrolyzate with a different DE in the range of about 2 to 9 to demonstrate that my conclusion in the previous declaration remains same at least when liquefied starch with a DE of less than 10 is used as a substrate.
 - 8. Formation of AA2G by the enzymes
 Experiment 8-1: Materials

Two kinds of partial starch hydrolyzates, "PINEDEX® #1" (DE 8±1) and "PINEDEX® #100" (DE 2-5), were purchased from Matsutani Chemical Industries Co., Ltd., Hyogo, Japan. (DEs of "PINEDEX® #1" and "PINEDEX® #100" are shown in the pamphlet of "PINEDEX®" distributed by Matsutani Chemical Industries Co., Ltd., Hyogo, Japan, copy attached.) A liquefied corn starch solution (DE 4.1) was prepared in our laboratory according to the following method. A corn starch was prepared into about 20% starch suspension, admixed with calcium carbonate to give a final concentration of 0.1%, adjusted to pH 6.5, and admixed with 0.3%/g-starch of "THERMAMYL® 60 L", an α -amylase commercialized by Novozymes Japan, Chiba, Japan, and then heated at 95°C for 15 minutes. After autoclaving at 120°C for 20 minutes, the reaction mixture was cooled to 53°C to obtain a liquefied corn starch solution. The DE of the liquefied corn starch was determined to be 4.1 by conventional Lane-Eynon method.

IMG from Arthrobacter globiformis A19 (FERM BP-7590) and RIAGase were prepared and partially purified, respectively, according to the method in Experiments 7 and 8 described in my previous declaration on November 28, 2008.

Experiment 8-2: Enzyme reaction

Three kinds of aqueous solution containing 5% (w/v) of L-ascorbic acid and 5% (w/v) of "PINEDEX® #1", "PINEDEX® #100", or the above liquefied corn starch, as a glucosyl donor, adjusted to pH 5.3 were used as substrates for IMG and RIAGase. To each substrate solution, 20 units/g-glucosyl donor of the partially purified IMG or 20 units/g-glucosyl donor of the partially purified RIAGase was added and subjected to an enzyme reaction at 50°C for 24 hours. After the reaction, the reaction mixture was boiled for 10 min to inactivate IMG or RIAGase.

Then, each reaction mixture, obtained by allowing IMG or RIAGase to act on the substrate, was admixed with 40 units/g-glucosyl donor of

plucoamylase commercialized by Seikagaku Corporation, Tokyo, Japan, to hydrolyze the remaining partial starch hydrolyzate into glucose and subjected to an enzyme reaction at 40°C for 17 hours. After the reaction, each reaction mixture was boiled for 10 min to inactivate glucoamylase. The resulting each solution was subjected to high performance liquid chromatography (HPLC) for determining the contents of AA2G, AA5G, and AA6G. HPLC analysis was carried out according to the method described in Experiments 5 and 7 of the specification of the present invention.

9. Experimental results

The results of Experiment 8 are summarized in Table 1.

Table 1

		Table		n reaction	mixture
		Dextrose	Content in reaction mixture (%, on a dry solid basis)		
Enzyme	Glucosyl	Equivalent			
	donor	(DE)	AA2G	AA5G	AA6G
		8±1	25.1	ND***	ND***
IMG*	PINEDEX® #1**	<u> </u>			
	PINEDEX®	2-5	25.6	ND***	ND***
	#100**				
	Liquefied corn	4.1	26.2	ND***	ND***
	starch		3.4.0	1.9	0.3
RIAGase	PINEDEX® #1**	8±1	14.8	1.3	
	PINEDEX®	2-5	14.9	2.0	0.3
	#100**				
	Liquefied corn	4.1	15.1	2.1	0.3
	starch	ater alobifo	770 (1	FERM BP-75	30)

^{*:} IMG from Arthrobacter globiformis A19 (FERM BP-7590)

***: Not detected

As evident from the results in Table 1, AA2G content in the reaction mixtures reached about 25.1 to 26.2%, on a dry solid basis, when IMG from

^{**:} Partial starch hydrolyzate commercialized by Matsutani Chemical Industries Co., Ltd., Hyogo, Japan

Arthrobacter globiformis A19 (FERM BP-7590) was allowed to act on "PINEDEX® #1" with a DE of 8±1, "PINEDEX® #100" with a DE of 2 to 5, and liquefied corn starch with a DE of 4.1, respectively. In addition, AA5G and AA6G, as by-products, were not detected in the reaction mixtures.

On the contrary to this, when RIAGase was allowed to act on the substrates with a DE of 8±1, 2 to 5, and 4.1, the AA2G contents in the reaction mixture were 14.8 to 15.1%, on a dry solid basis. Further, RIAGase produced AA5G and AA6G, as by-products, in amounts of about 2.0% and 0.3%, respectively.

10. Conclusion:

The above experimental results indicate that IMG from Arthrobacter globiformis A19 (FERM BP-7590) is superior to RIAGase in the productivity of AA2G when at least the DE of the substrate, i.e. liquefied starch, is in the range of about 2 to 9. Further, the above results indicate that IMG from Arthrobacter globiformis A19 (FERM BP-7590) is more suitable than RIAGase for producing AA2G at least the DE of the substrate, i.e. liquefied starch, is in the range of about 2 to 9, since IMG does not produce AA5G and AA6G as by products, while RIAGase does. It is believed that IMG from Arthrobacter globiformis A19 (FERM BP-7590) is a significantly different enzyme from RIAGase in the productivities of both AA2G and by products such as AA5G and AA6G.

I hereby further declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under 18 U.S.C. 1001,

and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Formyaki Niskimoto

NAME: Tomoyuki NISHIMOTO

26th day of August, 2010

DATE: 26th day of August, 2010

Attachment A

CURRICULUM VITAE

Name: Tomoyuki NISHIMOTO

Affiliation: Hayashibara Biochemical Laboratories, Inc.,

675-1, Fujisaki, Okayama-shi, Okayama,

Japan 702-8006

Tel: +81 86 276 8670

Date of Birth: December 26, 1961

Education: Granted and received a bachelor from Osaka

Prefecture University, Agricultural Department

in 1985.

Granted and received a master's degree from Osaka Prefecture University, Agricultural Department

in 1987.

Received a doctorate of Agriculture at Osaka

Prefecture University in 1998.

Brief Chronology of Employment:

1987 (April) Researcher, Hayashibara Co., Ltd.

1987 (July) Researcher, Toyama Medical and Pharmaceutical

University, under the employment of Hayashibara

Biochemical Laboratories, Inc.

1990 Researcher, Amase Institute, Research Center,

Hayashibara Biochemical Laboratories, Inc.

2002 Senior Scientist, Amase Institute, Research Center,

Hayashibara Biochemical Laboratories, Inc.

2004 Chief Scientist, Amase Institute,

Research Center, Hayashibara Biochemical

Laboratories, Inc.

2006 Chief Scientist, Glycoscience Institute,

Research Center, Hayashibara Biochemical

Laboratories, Inc.

2009 (April) Director, Glycoscience Institute,

Research Center, Hayashibara Biochemical

Laboratories, Inc.

Public Employment:

2006 (September) - Member of Editorial Board of Journal Applied

Glycoscience

2010 (April) - Affiliate associate professor of Hiroshima

University

2010 (April) - Part-time assistant professor of Kurashiki Sakuyo

University

List of Literatures

- 1. Tetsuya Nakada, Shoji Ikegami, <u>Tomoyuki Nishimoto</u>, Hiroto Chaen, Toshiyuki Sugimoto, and Masashi Kurimoto, <u>Purification and Characterization of Trehalase from Bacillus</u> sp. T3, Oyo Toshitsu Kagaku, 42, 231-236, 1995
- 2. <u>Tomoyuki Nishimoto</u>, Masayuki Nakano, Shoji Ikegami, Hiroto Chaen, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Existence of a Novel Enzyme Converting Maltose into Trehalose**, *Biosci. Biotech. Biochem.*, **59**, 2189-2190, 1995 *Short Communication*
- Tomoyuki Nishimoto, Masayuki Nakano, Tetsuya Nakada, Hiroto Chaen, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Purification and Properties of a Novel Enzyme, Trehalose Synthase, from** *Pimelobacter sp.* **R48**, *Biosci Biotech. Biochem.*, **60**, 640-644, 1996
- 4. <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Hiroto Chaen, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Purification and Characterization of a Thermostable Trehalose Synthase from** *Thermus aquaticus*, *Biosci. Biotech. Biochem.*, **60**, 835-839, 1996
- 5. Hiroto Chaen, Kazuhiko Maruta, Tetsuya Nakada, <u>Tomoyuki Nishimoto</u>, Takashi Shibuya, Michio Kubota, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Two Systems for Trehalose Biosynthesis in Bacteria**, *J. Appl. Glycosci.*, 43, 213-221, 1996 (in Japanese)
- Keiji Tsusaki, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Michio Kubota, Hiroto Chaen, Toshiyuki Sugimoto, Masashi Kurimoto, <u>Cloning and Sequencing of Trehalose</u> Synthase Gene from *Pimelobacter* sp. R48, *Biochem.Biophys Acta.*, 1290, 1-3, 1996 Short Sequence-Paper
- 7. <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Hiroto Chaen, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Action of a Thermostable Trehalose Synthase from** *Thermus aquaticus* **on Sucrose**, *Biosci. Biotech. Biochem.*, **61**, 898-899, 1997 *Note*
- Keiji Tsusaki, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Michio Kubota, Hiroto Chaen, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, Cloning and sequencing of trehalose synthase gene from *Thermus aquaticus* ATCC33923, *Biochem.Biophys Acta.*, 1334, 28-32, 1997 Short Sequence-Paper
- 9. Masashi Kurimoto, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Hiroto Chaen, Shigeharu Fukuda, and Yoshio Tsujisaka, **Synthesis by an** α-**Glucosidase of Glycosyl-trehaloses with an Isomaltosyl Residue**, *Biosci. Biotech. Biochem.*, **61**, 699-703, 1997
- 10. Hiroto Chaen, <u>Tomoyuki Nishimoto</u>, Takuo Yamamoto, Tetsuya Nakada, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, Formation of a Nonreducing Trisaccharide, Selaginose, from Trehalose by a Cell-free System of *Thermoanaerobium brockii*, *J. Appl. Glycosci.*, 46, 129-134, 1999

- Hiroto Chaen, Tetsuya Nakada, <u>Tomoyuki Nishimoto</u>, Nobue Kuroda, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, <u>Purification and Characterization of Thermostable Trehalose Phosphorylase from Thermoanaerobium brockii</u>, J. Appl. Glycosci., 46, 399-405, 1999
- Hiroto Chaen, Takuo Yamamoto, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, **Purification and Characterization of a Novel Phosphorylase, Kojibiose Phosphorylase, from** *Thermoanaerobium brockii*, *J. Appl. Glycosci.*, **46**, 423-429, 1999
- Hiroto Chaen, Tetsuya Nakada, Naoko Mukai, <u>Tomoyuki Nishimoto</u>, Shigeharu Fukuda, Toshiyuki Sugimoto, Masashi Kurimoto, and Yoshio Tsujisaka, Efficient Enzymatic Synthesis of Disaccharide, α-D-Galactosyl α-D-Glucoside, by Trehalose Phosphorylase from *Thermoanaerobacter brockii*, J. Appl. Glycosci., 48, 135-137, 2001 Note
- Hiroto Chaen, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, <u>Enzymatic Synthesis of Novel oligosaccharides from L-Sorbose</u>, <u>Maltose</u>, and <u>Sucrose Using Kojibiose Phosphorylase</u>, *J. Biosci. Bioengi.*, 92, 173-176, 2001
- Hiroto Chaen, <u>Tomoyuki Nishimoto</u>, Tetsuya Nakada, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, **Enzymatic Synthesis of Kojioligosaccharides Using Kojibiose Phosphorylase**, *J. Biosci. Bioengi.*, **92**, 177-182, 2001
- 16. Tomoyuki Nishimoto, Hajime Aga, Kazuhisa Mukai, Takaharu Hashimoto, Hikaru Watanabe, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, Purification and Characterization of Glucosyltransferase and Glucanotransferase Involved in the Production of Cyclic Tetrasaccharide in *Bacillus globisporus* C11, *Biosci. Biotecnol. Biochem.*, 66, 1806-1818, 2002
- 17. Tomoyuki Nishimoto, The Current Study of Cyclo-tetrasaccharide Focused on the Synthesizing System from Starch, Trends in Glycoscience and Glycotechnology, 14, 321-330, 2002 MINI REVIEW
- 18. Hajime Aga, Takanobu Higashiyama, Hikaru Watanabe, Tomohiko Sonoda, <u>Tomoyuki</u>
 <u>Nishimoto</u>, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka
 Production of Cyclic Tetrasaccharide from Starch Using a Novel Enzyme System
 from *Bacillus globisporus* C11, *J. Biosci. Bioengi.*, 94, 336-342, 2002
- Hajime Aga, <u>Tomoyuki Nishimoto</u>, Mieko Kuniyoshi, Kazuhiko Maruta, Hiroshi Yamashita, Takanobu Higashiyama, Tetsuya Nakada, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, 6-α-Glucosyltransferase and 3-α-Isomaltosyltransferase from *Bacillus globisporus* N75, *J. Biosci. Bioengi.*, 95, 215-224, 2003
- 20. Hideki Okada, Eri Fukushi, Shuichi Onodera, <u>Tomoyuki Nishimoto</u>, Jun Kawabata, Masanori Kikuchi, Norio Shiomi, **Synthesis and structural analysis of five novel**Oligosaccharides prepared by glucosyltransfer from β-D-glucose 1-phosphate to isokestose and nystose using *Thermoanaerobacter brockii* kojibiose phosphorylase,

- Carbohyd. Res., 338, 879-885, 2003
- Tomoyuki Nishimoto, Hajime Aga, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, **Production of Cyclic Tetrasaccharide with**6-α-Glucosyltransferase and α-Isomaltosyltransferase, J. Appl. Glycosci., **51**, 135-140, 2004
- Hikaru Watanabe, Masayuki Nakano, Kazuyuki Oku, Hajime Aga, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, **Cyclic**Tetrasaccharide in *Sake* Lees, *J. Appl. Glycosci.*, **51**, 345-347, 2004
- Takanobu Higashiyama, Hikaru Watanabe, Hajime Aga, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, <u>Enzymatic</u> Synthesis of a β-D-galactopyranosyl cyclic tetrasaccharide by β-galactosidases, *Carbohyd. Res.*, 339, 1603-1608, 2004
- Hajime Aga, Takanobu Higashiyama, Hikaru Watanabe, Tomohiko Sonoda, Ritsuko Yuen, Tomoyuki Nishimoto, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, Enzymatic Synthesis of Glycosyl Cyclic Tetrasaccharide with 6-α-Glucosyltransferase and 3-α-Isomaltosyltransferase, J. Biosci. Bioeng., 98, 287-292, 2004
- Takuo Yamamoto, Kazuhiko Maruta, Kazuhisa Mukai, Hiroshi Yamashita, Tomoyuki Nishimoto, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, Cloning and Sequencing of Kojibiose Phoshorylase Gene from Thermoanaerobacter brockii ATCC35047, J. Biosci. Bioeng., 98, 99-106, 2004
- Tomoyuki Nishimoto, Novel Process for Producing Cyclic Tetrasaccharide and Functions of the Cyclic Tetrasaccharide, Nippon Nogekagaku Kaishi, 78, 866-869, 2004
 MINI REVIEW (in Japanese)
- Hikaru Watanabe, Takanobu Higashiyama, Hajime Aga, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, **Enzymatic** synthesis of a 2-O-α-D-glucopyranosyl cyclic tetrasaccharide by Kojibiose Phoshorylase, Carbohyd. Res., 340, 449-454, 2005
- Takuo Yamamoto, Kazuhisa Mukai, Kazuhiko Maruta, Hikaru Watanabe, Hiroshi Yamashita, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Hiroto Chaen, and Shigeharu Fukuda, Hyper Expression of Kojibiose Phoshorylase Gene and Trehalose Phosphorylase Gene from *Thermoanaerobacter brockii* ATCC35047 in *Bacillus subtilis* and Selaginose Synthesis Utilizing Two Phosphorylases, *J. Biosci. Bioeng.*, 100, 343-346, 2005
- Tetsuya Nakada, Masashi Kurimoto, <u>Tomoyuki Nishimoto</u>, Torajiro Nakahara, Shoji Ikegami, Hiroto Chaen, Shigeharu Fukuda, and Toshiyuki Sugimoto, **Purification and Some Properties of β-Fructofuranosidase from** *Bacillus* sp. **V230**, *ITE Letters on Batteries, New Technologies & Medicine*, Vol. 6, No. 2, 145-150, 2005
- 30. Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Hajime Aga, Michio Kubota, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, <u>Enzymatic Synthesis of a Novel cyclic</u>

- Pentasaccharide consisting of α-D-Glucopyranose with 6-α-Glucosyltransferase and 3-α-Isomaltosyltransferase, Carbohyd. Res., 340, 1577-1582, 2005
- Kazuhisa Mukai, Hikaru Watanabe, Kazuyuki Oku, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Hiroto Chaen, Shigeharu Fukuda and Masashi Kurimoto, **An Enzymatically Produced Novel cyclic Trtrasaccharide**, *cyclo*-{→6}-α-D-Glcp-(1→4)-α-D-Glcp-(1→6)-α-D-Glcp-(1→4)-α-D-Glcp-(1→4) (cyclic maltosyl-(1→6)-maltose), from Starch, *Carbohydr. Res.*, 340, 1469-1474, 2005
- 32. <u>Tomoyuki Nishimoto</u>, Study of Trehalose-relating Enzymes, J. Appl. Glycosci., 53, 57-64, 2006 Proceedings
- Kazuhiko Maruta, Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Hiroto Chaen, Shigeharu Fukuda, Masashi Kurimoto, and Yoshio Tsujisaka, Acceptor Specificity of Trehalose Phosphorylase from *Thermoanaerobacter brockii*: Production of Novel Nonreducing Trisaccharide, 6-O-α-D-Galactopyranosyl Trehalose, J. Biosci. Bioeng., 101, 385-390, 2006
- Takuo Yamamoto, <u>Tomoyuki Nishimoto</u>, Hiroto Chaen, and Shigeharu Fukuda, Improvement of the Enzymatic Properties of Kojibiose Phoshorylase from Thermoanaerobacter brockii by Random Mutagenesis and Chimerization, J. Appl. Glycosci., 53, 123-129, 2006 Symposium proceedings
- Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Tomohiko Sonoda, Michio Kubota, Hiroto Chaen and Shigeharu Fukuda, An Enzymatically Produced Novel Cyclomaltopentaose Cyclized from Amylose by an α -(1 \rightarrow 6)-linkage, cyclo-{ \rightarrow 6)- α -D-Glcp-(1 \rightarrow 4)- α -D-Glcp-(1 \rightarrow 4)-
- Res., **341**, 957-963, 2006
- Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Kazuhisa Mukai, Michio Kubota, Hiroto Chaen, And Shigeharu Fukuda, A Novel Glucanotransferase from *Bacillus circulans* Strain That Produces a Cyclomaltopentaose cyclized by an α-(1→6)-linkage, *Biosci. Biotecnol. Biochem.*, 70, 1954-1960, 2006
- Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Hiroto Chaen, and Shigeharu Fukuda, Cloning, Sequencing, and Expression of the Genes Encoding an Isocyclomaltooligosaccharide Glucanotranferase and an α-Amylase from a *Bacillus circulans* Strain, *Biosci. Biotecnol. Biochem.*, 70, 2690-2702, 2006
- Hikaru Watanabe, Rohko Takakura-Yamamoto, Mayumi Kurose, Kenshi Yoshida, Kazuyuki Oku, Ikuo Sawatani, <u>Tomoyuki Nishimoto</u>, Michio Kubota, Hiroto Chaen, and Shigeharu Fukuda, <u>Production of Isocyclomaltopentaose from Starch Using Isocyclomaltooligosaccharide Glucanotransferase</u>, *Biosci. Biotecnol. Biochem.*, 70, 3013-3018, 2006
- Takuo Yamamoto, Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Hajime Aga, Michio Kubota, Hiroto Chaen, and Shigeharu Fukuda, <u>Acceptor Recognition of Kojibiose Phoshorylase from Thermoanaerobacter brockii: Syntheses of Glycosyl Glycerol and myo-Inositol, J. Biosci. Bioeng.</u>, 101, 427-433, 2006

- Kazuhiko Maruta, Michio Kubota, Hiroshi Yamashita, <u>Tomoyuki Nishimoto</u>, Hiroto Chaen and Shigeharu Fukuda, Creation of a Novel Hydrolase by Site-directed Mutagenesis of Malto-oligosyltrehalose Synthase, *J. Appl. Glycosci.*, **53**, 199-203, 2006
- Tomoyuki Nishimoto, Kazuyuki Oku, and Kazuhisa Mukai, Two Systems for Producing Cyclic Tetrasaccharide Using Starch as a Substrate, Kagaku To Seibutsu, 44, 539-550, 2006 (in Japanese)
- Takaharu Hashimoto, Mayumi Kurose, Kazuyuki Oku, <u>Tomoyuki Nishimoto</u>, Hiroto Chaen, Shigeharu Fukuda, and Yoshio Tsujisaka, <u>Digestibility and Suppressive Effect on Rats' Body Fat Accumulation of Cyclic Tetrasaccharide</u>, *J. Appl. Glycosci.*, **53**, 233-239, 2006
- Watanabe H, Nishimoto T, Chaen H, Fukuda S, A Novel Glucanotransferase that Produces a Cyclomaltopentaose Cyclized by an α-1,6-Linkage, J. Appl. Glycosci., 54, 109-118, 2007 Symposium proceedings
- Hayakawa N, Kigawa R, <u>Nishimoto T</u>, Kakamoto K, Fukuda S, Kimishima T, Oka Y and Kawanobe W, Characterization of Furunori (Aged Paste) and Preparation of a Polysaccharide Similar to Furunori, *Studies in Conservation*, 52, No.3, 2007
- Takahashi N, Fukushi E, Onodera S, Benkeblia N, <u>Nishimoto T</u>, Kawabata J, Shiomi N, Three novel oligosaccharides synthesized using *Thermoanaerobacter brockii* kojibiose Phosphorylase, *Chemistry Central Journal*, 1, 18, 2007
- Tetsuya Mori, <u>Tomoyuki Nishimoto</u>, Takanori Okura, Hiroto Chaen, and Shigeharu Fukuda, <u>Purification and Characterization of Cyclic Maltosyl-(1→6)-Maltose Hydrolase and α-Glucosidase from an *Arthrobacter globiformis* Strain, *Biosci. Biotecnol. Biochem.*, 72, 1673-1681, 2008</u>
- Motohiro Shizuma, Taro Kiso, Hisashi Terauchi, Yoshio Takai, Hitoshi Yamada, <u>Tomoyuki Nishimoto</u>, Daisuke Ono, Osamu Shimomura, Ryoki Nomura, Yoshikatsu Miwa, Masaki Nakamura, and Hirofumi Nakano, Evaluation of Chiral Amino Acid Discrimination by a Permethylated Cyclic Tetrasaccharide, cyclo-{→6}α-D-Glcp-(1→3)-α-D-Glcp-(1→6)-α-D-Glcp-(1→3)-α-D-Glcp-(1→}, Using FAB MASS Spectrometry, Chemistry Letters, 37, 1054-1055, 2008
- Tetsuya Mori, <u>Tomoyuki Nishimoto</u>, Kazuhisa Mukai, Hikaru Watanabe, Takanori Okura, Hiroto Chaen, and Shigeharu Fukuda, <u>Enzymes Involved in the Biosynthesis and Degradation of Cyclic Maltosyl-Maltose in *Arthrobacter globiformis* M6, *J. Appl. Glycosci.*, **56**, 127-136, 2009</u>
- Keiji Tsusaki, Hikaru Watanabe, <u>Tomoyuki Nishimoto</u>, Takuo Yamamoto, Michio Kubota, Hiroto Chaen, and Shigeharu Fukuda, **Structure of a Novel Highly branched** α-**Glucan enzymatically produced from Maltodextrin**, *Carbohydrate Research*, **344**, 2151-2156, 2009